

WRF-Chem V3.4.1: A Quick Review Of How To Set-Up & Run

Steven Peckham

WRF-Chem


- It is assumed that the user of WRF-Chem :
 - is *very familiar* with the WRF model system
 - have run WPS
 - and has made more than one weather simulation using WRFV3
 - know FORTRAN and C and can edit code, recompile, etc.
- The chemistry code is available from WRF web page.
 - Questions: Send email to WRF-Chem help (wrfchemhelp.gsd@noaa.gov)
 - Web page: www.wrf-model.org/WG11
- Test data is available as well (tutorial exercises)
 - Small domain (41x41x31 grid points, 100 km horiz. spacing)

WRF-Chem

- Compile WRF-Chem code (already done for tutorial)
 - Set environmental variables
 - Define which model core to build (use ARW only).
 - `setenv WRF_EM_CORE 1`
 - `setenv WRF_NMM_CORE 0`
 - Chemistry code is to be included in the WRF model build
 - `setenv WRF_CHEM 1`
 - Kinetic Pre-Processor (KPP) code
 - `setenv WRF_KPP 1` => if KPP is to be included
 - `setenv WRF_KPP 0` => if KPP is NOT to be included
 - `setenv FLEX_LIB_DIR /usr/lib`
 - `setenv YACC '/usr/bin/yacc -d'`
 - Configure and issue “`compile em_real`” command
 - Save compile output to file
 - Check results for errors and check known problems web page if no `wrf.exe`

WRF-Chem Emissions

WRF-ARW Modeling System Flow Chart



WRF-Chem Emissions

- Two sources of anthropogenic emissions available on ftp site:
 - RETRO (.5 degree, monthly) and EDGAR (10 degree, annual)
 - Run Prep_sources_chem (Tutorial exercise 2)
 - Surface data only (kemit=1)
 - National Emissions Inventory (NEI-2005) for U.S.
 - Point source and stack plume rise emissions computed offline and applied to 3d domain (kemit \geq 1)
 - Run without chemistry first as mean wind profile is needed!
 - Both include programs to map to WRF grid; binary output files

WRF-Chem Emissions


- Can use other external emissions data
 - Start with “raw” emissions data
 - Specify the speciation for the desired chemical mechanism
 - Prepared the 3-D (or 2-D) anthropogenic emissions data set
 - Map data onto your WRF-Chem simulation domain
 - Output data
 - Goal: have data in a WRF data file to run with model

WRF-Chem Emissions

- The available methodology for emissions uses a convert program
 - Program called `convert_emiss.exe` (compile `emi_conv`)
 - Reads header information from a WRF input file
 - Reads binary emissions data
 - Writes a WRF netCDF data file
- `convert_emiss` is very simple. Uses just a few namelist settings.
 - `emiss_opt=3` – NEI emissions for U.S.A.
 - `emiss_opt=5` – RETRO/EDGAR global emission
- Data is read in via `auxinput5` when running `wrf.exe`
 - `auxinput5_inname` = 'wrfchemi_<hr>z_d<domain>', (optional)
 - `io_form_auxinput5` = 2,
- Chpt. 3 and Appendix B of User's Guide for more information
- Users can create input data files through any other methodology

WRF-Chem Biogenic Emissions

WRF-ARW Modeling System Flow Chart



WRF-Chem

Biogenic Emissions

- 4 choices for Biogenic emissions
- Option 1: No biogenic emissions ($\text{bio_emiss_opt} = 0$):
 - Provide biogenic emissions through anthropogenic input.
 - No additional input data files.
- Option 2 ($\text{bio_emiss_opt} = 1$): (good default option)
 - Landuse based emissions following Guenther et al (1993, 1994), Simpson et al. (1995). Emissions depends on both temperature and photosynthetic active radiation.
 - No additional input data files.
 - Small number of vegetation types (errors?)

WRF-Chem

Biogenic Emissions

- Option 3 (bio_emiss_opt = 2):
 - User specified from external data source
 - Biogenic Emissions Inventory System (BEIS) version 3.14 [*Vukovich and Pierce, 2002*] with land-use obtained from the Biogenic Emissions Landuse Database version 3 (BELD3) [*Pierce et al., 1998*].
 - Static 2-D surface data provided in input data file and are modified according to the environment
 - Data is read in via auxinput6 when running real.exe
 - auxinput6_inname = 'wrfbiochemi_d01',
 - io_form_auxinput6 = 2,

WRF-Chem

Biogenic Emissions

- Option 4 (bio_emiss_opt = 3): MEGAN (best choice?!)
 - Separate program made available by NCAR/ACD
 - Global data with base resolution of ~ 1 km
 - Leaf Area Index, vegetation type, emission factors
 - Steps:
 1. Download MEGAN code from NCAR/ACD
 - megan_bio_emiss.tar
 - megan.data.tar.(when uncompressed ~ 28 GB)

<http://acd.ucar.ued/~guenter/MEGAN/MEGAN.html>

WRF-Chem

Biogenic Emissions

- Option 4 (bio_emiss_opt = 3): MEGAN
 - Steps:
 2. Compile megan_bio_emiss
 3. Create wrfbiochemi_d01 data file using:
 - wrfinput,
 - RAW MEGAN data files,
 - settings in megan_bio_emiss.input file
 - About 10 Gb of memory

WRF-Chem

Biogenic Emissions

- Option 4 (bio_emiss_opt = 3): MEGAN
 - Steps:
 4. View wrfbiochemi_d01 data file to verify data is correct
 5. Run real.exe and wrf.exe
 - Add ne_area setting to the WRF chemistry namelist!!!
 - » ne_area = number of chemical species in chem_opt

<http://acd.ucar.ued/~guenter/MEGAN/MEGAN.html>

WRF-Chem

Biomass Burning Emissions

- 2 choices for biomass burning emissions
- Option 1: No biomass emissions (biomass_burn_opt = 0):
 - No additional input data files.
- Option 2 (biomass_burn_opt = 1):
 - Use prep_chem_sources program to read WFABBA, or MODIS data
 - Convert binary data to wrffirechemi_d01 input file
 - Use FINN historical data
 - Data read in through auxinput7 when running real.exe

WRF-Chem

Dust Emissions

- 3 choices for dust emissions
- Option 1: No dust emissions (`dust_opt = 0`):
 - No additional input data files.
- Option 2 (`dust_opt = 1`):
 - Need to include surface erosion data in WPS
 - Use new GEOGRID table for running `geogrid.exe`
 - Dust data included in `wrfinput` file
- Option 3 (`dust_opt = 3`)
 - AFWA scheme uses same method as option 2
- Work tutorial exercise 1 for more information.


WRF-Chem

GOCART Background Data

- Includes DMS as well as GOCART
 - From running prep_chem_sources with GOCART included
 - Planned to be moved to WPS one of these days
- Run prep_chem_sources program to produce external binary data files
- Convert binary data files to WRF input files
 - chem_opt = 300 or 301 and/or dmsemis_opt=1
- Data read by real.exe through auxinput8
 - File name wrfchemi_gocart_bg_d01

WRF-Chem Initial Conditions

WRF-ARW Modeling System Flow Chart




WRF-Chem Chemical I.C.s

- WPS gives meteorology I.C.s
- Problem: Need 3-D state of atmospheric chemistry for model initialization
 - Every 12 hours balloons used to measure state of atmosphere
 - Instruments rarely measure chemistry
 - Temperature, relative humidity, wind speed and direction
 - Occasionally measure ozone
 - No other consistent measurements of chemistry
 - Satellites measure AOD
 - Vertical integral
 - Distinguish between type (carbon, sulfate, etc.) aerosols?
- Solution: WRF-Chem model provides chemical I.C.s

WRF-Chem Chemical I.C.s

- Option 1 – Idealized profile
 - Every grid point gets same chemical profile
 - Assumes August in North America for I.C.
 - Set `chem_in_opt = 0` in namelist, run `real.exe`
- Option 2 – WRF/Chem Forecast
 - Use previous forecast for chemistry I.C.
 - Set `chem_in_opt = 1` in namelist, run `real.exe`
- Option 3 – Use other model data for I.C.
 - User needs to modify the `wrfinput` file after running `real.exe`
 - No `chem_in_opt` setting necessary (`chem_in_opt=0` default)

WRF-Chem Boundary Conditions



WRF-Chem Chemistry B.C.s

- External tools under development to provide global model data as BC and initial conditions
- Older program available: wrfchembc (Rainer Schmitz - Univ. of Chile)
 - Available code runs with MPI-MATCH & RAQMS data
 - Adds lateral boundary data for chemical species to wrfbdy_d01
 - User specifies which chemical species to use
 - Need to choose chemical species from global model
 - Need to speciate global model data for WRF-Chem chemistry
 - Requires knowledge from user regarding chemistry (not turn-key)
- wrfinput_d01 not modified
 - Can result in differences near boundaries at start of simulation

WRF-Chem Chemistry B.C.s

- Other groups are exploring other possible ways to generate input/B.C. data for WRF-Chem
 - NCAR/ACD has a program available if using MOZART
- MOZBC sets space and time-varying chemical initial (IC) and boundary conditions (BC)
 - global model output (MOZART-4 or CAM-Chem)

MOZBC : <http://www.acd.ucar.edu/wrf-chem/download.shtml>


MOZART data (2004-2008):

<http://www.acd.ucar.edu/wrf-chem/mozart.shtml>

- Note: MOZART/CAM-Chem data are interpolated only in space.

WRF-Chem Chemistry B.C.s

- Program will fill the chemical fields in your `wrfinput_d<nn>` and `wrfbdy_d<nn>` files with global model output.




- To enable chemical IC and BC when running WRF-Chem set in `namelist.input`:
have_bcs_chem = .true.

WRF-Chem Chemistry B.C.s

- What if you have different GCM data?
- Methodology is the same
 - Read global model chemistry data
 - Skip over if not a desired chemistry species
 - Determine grid point location on WRF-Chem grid
 - If at boundary, interpolate data to WRF-Chem grid
 - Speciate to your chemical mechanism
 - Once completed reading/interpolating/speciating global data:
 - Open wrfbdy_d01 data file
 - Write boundary data to wrfbdy_d01

WRF-Chem Namelist

WRF-ARW Modeling System Flow Chart



WRF-Chem Namelist

- **Time control namelist options**
- A few of the chemistry namelist options
 - More details provided in Chapter 4 of User's Guide

WRF-Chem Namelist

- Time control namelist
 - Chemistry input fields come in through auxiliary input ports
 - For example, biogenic emissions use auxinput 6

&time_control

...

auxinput6_inname	= 'wrfbiochemi_d<domain>,'
auxinput6_interval_m	= 1440,
io_form_auxinput6	= 2,

Recall:

Defining a variable-set for an I/O stream

- **Fields are added to a variable-set on an I/O stream at compile-time with Registry**

#	Type	Sym	Dims	Use	Tlev	Stag	IO	Dname	Descrip
state	real	u	ikjb	dyn_em	2	X	i01rhusdf	"U"	"X WIND COMPONENT"

IO is a string that specifies if the variable is to be subject to initial, restart, or history I/O. The string may consist of '**h**' (subject to history I/O), '**i**' (initial dataset), or '**r**' (restart dataset). The 'h', 'r', and 'i' specifiers may appear in any order or combination.

The 'h' and 'i' specifiers may be followed by an optional integer string consisting of '0', '1', ..., '9'. Zero denotes that the variable is part of the principal input or history I/O stream. The characters '1' through '9' denote one of the auxiliary input or history I/O streams.

WRF-Chem Registry

- Thus, in registry.chem

```

state real - i+jf emis_ant - - - - "Anthropogenic Emissions" ""
state real e_iso i+jf emis_ant 1 Z i5r "E_ISO" "Isoprene EMISSIONS" "mol km^-2 hr^-1"
state real e_so2 i+jf emis_ant 1 Z i5r "E_SO2" "EMISSIONS" "mol km^-2 hr^-1"
state real e_no i+jf emis_ant 1 Z i5r "E_NO" "EMISSIONS" "mol km^-2 hr^-1"

#
state real e_bio ijo misc 1 Z r "E_BIO" "EMISSIONS" "ppm m/min"
state real sebio_iso ij misc 1 - i6r "sebio_iso" "Reference biog emiss" "mol km^-2 hr^-1"
state real sebio_oli ij misc 1 - i6r "sebio_oli" "Reference biog emiss" "mol km^-2 hr^-1"

# additional arrays needed for biomass burning emissions input
state real - iljf ebu_in - - - - "Biomass burnung input" ""
state real ebu_in_no iljf ebu_in 1 - i{7} "ebu_in_no" "EMISSIONS" "mol km^-2 hr^-1"
state real ebu_in_co iljf ebu_in 1 - i{7} "ebu_in_co" "EMISSIONS" "mol km^-2 hr^-1"

# Input for GOCART: Background chemistry, erodible surface emissions map
state real backg_oh ikj misc 1 - i8r "BACKG_OH" "Background OH" "volume mixing ratio"
state real backg_h2o2 ikj misc 1 - i8r "BACKG_H2O2" "Background H2O2" "volume mixing ratio"

```


WRF-Chem Namelist

- For the chemistry variables to come in via auxiliary port
 - Registry set for input via auxiliary port

Auxiliary port number	Description
5	Anthropogenic emissions
6	Biogenic emissions
7	Surface biomass burning data
8	GOCART background fields
12	External chemistry fields (wrfout data from previous run)
13	Volcanic Ash emissions
14	Aircraft emissions
15	Green House Gas emissions

WRF-Chem Namelist

- For the chemistry variables to come in via auxiliary ports (cont.)
 - Namelist set in time_control

```
&time_control
...
auxinput6_inname = 'wrfbiochemi_d01',
auxinput7_inname = 'wrffirechemi_d<domain>',
auxinput8_inname = 'wrfchemi_gocart_bg_d<domain>',
auxinput12_inname = 'wrf_chem_input',
auxinput13_inname = 'wrfchemv_d<domain>',
auxinput5_interval_m = 86400, 86400, 60,
auxinput7_interval_m = 86400, 86400, 60,
auxinput8_interval_m = 86400, 86400, 60,
auxinput13_interval_m = 86400, 86400, 60,
io_form_auxinput2 = 2,
io_form_auxinput5 = 2,
io_form_auxinput6 = 0,
io_form_auxinput7 = 0,
io_form_auxinput8 = 0,
io_form_auxinput12 = 0,
io_form_auxinput13 = 0,
```


WRF-Chem Namelist

A few of the Chemistry namelist options

- More details provided in Chapter 4 of WRF-Chem User's Guide

WRF-Chem Namelist

- Chemistry control namelist

Chem_opt	Description
0	No chemistry
1 - 40	Chemical mechanisms (RADM2, CBMZ), tracer options (chem_opt=13 to 17)
101 - 200	Options covering RADM2, CBMZ, MOZART, SAPRC99, NMHC9 chemical mechanisms using KPP.
300 – 303	GOCART aerosol options
400 – 403	Dust and Volcano options (volcanic and surface lofted)

WRF-Chem Namelist

emiss_opt	Description
0	no anthropogenic emissions
2	RADM2 anthropogenic emissions
3	RAM2/MADE/SORGAM anthropogenic emissions
4	CBMZ/MOSAIC anthropogenic emissions
5	GOCART RACM_KPP emissions
6	GOCART simple emissions
7	MOZART emissions .
8	MOZCART (MOZART + GOCART aerosols) emissions
13	SAPRC99 emissions
16	CO2 tracer emissions
17	Green House Gas emissions

Remember: emiss_opt sets emissions structure (registry.chem)

#emission package definitions

```
package eradmsorg    emiss_opt==3
emis_ant:e_iso,e_so2,e_no,e_no2,e_co,e_eth,e_hc3,e_hc5,e_hc8,e_xyl,e_ol2,e_olt,e_oli,e_tol,e_csl,e_hch
o,e_ald,e_ket,e_or2,e_nh3,e_pm25i,e_pm25j,e_pm_10,e_eci,e_ecj,e_orgi,e_orgj,e_so4i,e_so4j,e_no3i,e_
no3j,e_naaj,e_naai,e_orgi_a,e_orgj_a,e_orgi_bb,e_orgj_bb
```

```
package ecptec      emiss_opt==5
emis_ant:e_iso,e_so2,e_no,e_no2,e_co,e_eth,e_hc3,e_hc5,e_hc8,e_xyl,e_ol2,e_olt,e_oli,e_tol,e_csl,e_hch
o,e_ald,e_ket,e_or2,e_nh3,e_pm_25,e_pm_10,e_oc,e_sulf,e_bc
```

Anthropogenic CO₂, CO and CH₄ emissions:

```
package eco2        emiss_opt==16    emis_ant:e_co2,e_co2tst,e_co
```

```
package eghg        emiss_opt==17    emis_ant:e_co2,e_co2tst,e_co,e_cotst,e_ch4,e_ch4tst
```


WRF-Chem Namelist

cu_rad_feedback	Description
.false.	No feedback from the parameterized convection to the atmospheric radiation and the photolysis schemes. (logical)
.true.	Feedback from the parameterized convection to the radiation schemes turned on. (logical) - use Grell cumulus scheme
progn	
0	Turns off prognostic cloud droplet number in the Lin et al. microphysics
1	Prognostic cloud droplet number included in the Lin et al. This effectively turns the Lin et al. scheme into a second-moment microphysical scheme. If set with chem._opt=0 a default prescribed aerosol concentration is used.

WRF-Chem Namelist

cldchem_onoff	Description
0	cloud chemistry turned off in the simulation, also see the “chem_opt” parameter
1	cloud chemistry turned on in the simulation, also see the “chem_opt” parameter
wetscav_onoff	
0	wet scavenging turned off in the simulation, also see the “chem_opt” parameter
1	wet scavenging turned on in the simulation, also see the “chem_opt” parameter

WRF-Chem Namelist

- Useful option for real time forecasts
- `chem_in_opt = 0`
 - No chemical initial analysis derived from observations
 - Default initial state for N. America summer
- `chem_in_opt=1`
 - Use forecast for initial chemical fields
 - Good choice for real time forecast runs
 - Works well as lower tropospheric air quality mostly depends on emissions
 - `real.exe` reads in forecast data through `auxinput12`

WRF-Chem Namelist


- `chem_in_opt=1`: additional methodology
 - Set namelist option `chem_in_opt = 1`
 - Update dates/times of simulation in `namelist.input` for your forecast
 - Copy or link `wrfout` file to a “`wrf_chem_input`” data file
 - Set `auxinput12` namelist options

```
ln -s $outdir/wrfout_d01_2007-06-15-12:00:00 wrf_chem_input_d01
```

- `Real.exe` gives a message indicates that model is being initialized with previous forecast
- Do tutorial exercise 5 to better understand methodology

WRF-Chem Simulation

WRF-ARW Modeling System Flow Chart



Running WRF-Chem

Option 1: Run without chemistry

- Get copy of WRF code in your home directory, compile (w/o chem)
- Get WPS into your home directory, compile
- Get initialization data (e.g., global model data)
- Build "met_em" input data files using WPS
- Set options in namelist.input
- Run real.exe
- Run wrf.exe
- Check results

Running WRF-Chem

Option 2: Run with dust only (tutorial exercise 1)

- Get copy of WRF-Chem code in your home directory

```
cp -R /wrfhelp/SOURCE_CODE/WRF-CHEM/WRFV3 WRFV3
```

The tutorial code is pre-compiled (skip section 1 in the quick start guide).

- Set WPS links to include dust
 - link GEOGRID.TBL.ARW_CHEM to GEOGRID.TBL
 - Should have EROD data
- Build "met_em" input data files (include dust (EROD) fields)
- Set options in namelist.input
- Run real.exe with the dust only chemistry option turned on.
(Save the wrfinput_d01 data file for use later on.)
- Run wrf.exe and check results

Running WRF-Chem

Option 3: Run with chemistry/aerosols (exercises 2 & 3)

- Start with WPS output and wrfinput file from option 2 then...
- Compile the prep_chem_sources program
- Set namelist input options (directories, on/off switches)
- Run the prep_chem_sources program
- Link output files to your meteorology run directory
- Retrieve the wrfinput file from your meteorology or dust only run

Running WRF-Chem

Option 3: Run with chemistry/aerosols (cont.)

- Set namelist options
 - Auxiliary input stream 5 for anthropogenic emissions
 - auxinput5_interval = 60
 - io_form_auxinput5 = 2
 - chem_opt = 301
 - emiss_opt=5
 - kemit=1
 - dust_opt=1
 - biomass_burn_opt=1
- Run convert_emiss.exe
 - Produces wrfchemi_d01, wrfchemi_gocart_backg_d01, wrffirechemi_d01 input data files.
 - No volcanic ash emissions as chem_opt and emiss_opt_vol not set.
- Copy emissions files to WRFV3/test/em_real run directory

Running WRF-Chem

Option 3: Run with chemistry/aerosols (cont.)

- Run real.exe to include chemistry
- If including fixed surface emissions (biogenic, fire, GOCART background, etc.):
 - Set namelist options to include additional inputs
 - Auxinput6 – biogenic emissions
 - Auxinput7 – biomass burning emissions
 - Auxinput8 – GOCART background fields
 - Auxinput12 – Include previous run's chemistry fields
- After running real.exe
 - Have wrfinput_d01 and wrfbdy files with chemistry fields
 - Should get messages showing chemistry is in run.
 - Verify data

Running WRF-Chem


Option 3: Run with chemistry/aerosols (cont.)

- Run wrf.exe to produce wrfout data files
 - Set namelist options to include additional inputs
 - Auxinput5 – anthropogenic emissions
 - Turn off namelist options that are included in wrfinput file
 - Auxinput7 – biomass burning emissions
 - Auxinput8 – GOCART background fields
 - Auxinput12 – Include previous run's chemistry fields
 - Should get messages regarding the reading of
 - Anthropogenic emissions
 - If you do not get these messages, an error is likely

After Running WRF-Chem

- And still other options (tracer, volcano, etc.)
- For any of the run time options
 - Check the text output
 - Make sure you are getting the messages you expect
 - Look for any warning/error messages
 - Check the model output (ncview)
 - Confirm that emissions data is being read into simulation
 - Error in kemit will result in no anthropogenic emissions data
 - Error in chem_opt, io_form_auxinput5?
 - Other namelist options correct so you have data?
 - » `Max<some array> = 1.e-16` is not data. It is noise.
 - Make plots of simulation results

WRF-Chem Visualization



WRF-Chem Visualization

- Your favorite netCDF data file viewer to examine results
 - ncview, ncbrowse, etc.
- Other standard WRF visualization tools work with the chemistry variables as well as the meteorology
 - ARWpost (NCL, VIS5D)
 - Grads
 - Etc.

