
Aerosol modeling with WRF/Chem

Jan Kazil

University of Colorado / NOAA Earth System Research Laboratory

WRF/Chem Tutorial, 22 July 2013

Part I - Introduction

- Overview of ...
 - Aerosol
 - Aerosol processes and life cycle
 - Model treatment of aerosol
 - WRF/Chem aerosol schemes

Part II – The details


- Representing the aerosol size distribution
- Walk through the WRF/Chem aerosol schemes
 - How they work and what they do
 - Coupling to other processes
 - ◆ Gas phase chemistry
 - ◆ Aqueous chemistry
 - ◆ ...
- Hint on how to tell WRF/Chem what to do
- Resources

Part I – Introduction

Aerosol




Aerosol life cycle and processes



Model treatment of aerosol

Individual processes or sets of processes are treated by “modules” or “schemes”.




WRF/Chem aerosol schemes

- **An efficient aerosol scheme from the GOCART model**
 - No size information for sulfate, BC, OC
 - Size information for dust and sea salt
 - No secondary organic aerosol (SOA)
- **Modal Aerosol Dynamics Model for Europe – MADE**
 - 3 log-normal modes
 - Inorganic, organic aerosol, SOA
- **Model for Simulating Aerosol Interactions and Chemistry (MOSAIC)**
 - Sectional model, 4 or 8 bins
 - Inorganic, organic aerosol, SOA
- **MAM – Modal Aerosol Model (new in WRF/Chem 3.5) from CAM5**
 - 3 or 7 log-normal modes
 - Inorganic, organic aerosol, SOA, sea salt, BC, mineral dust
- **Simple sectional (bin) scheme for volcanic ash aerosol**

Part II – The details

Bulk aerosol schemes

- Only total mass of aerosol compounds is known



- No information on
 - Particle number
 - Aerosol size distribution

Aerosol size distribution needs to be assumed for:


- radiative transfer
- response of cloud properties to aerosol number

- Numerically efficient
- Useful when focus is on complex gas phase chemistry

→ **GOCART (+ size resolved dust and sea salt)**


Modal aerosol schemes

Twin Otter data (black)



Modal aerosol schemes

Twin Otter data (black)



$$\frac{dN}{dD} = \frac{N}{\sqrt{2\pi \ln(\sigma)} D} e^{-\frac{1}{2} \left[\frac{\ln(D/\mu)}{\ln(\sigma)} \right]^2}$$


$$\frac{dN}{dD} \rightarrow N = 8195 \text{ cm}^{-3}$$
$$\mu = 18.22 \text{ nm}$$
$$\sigma = 1.42$$

$$\frac{dN}{dD} \rightarrow N = 12732 \text{ cm}^{-3}$$
$$\mu = 68.44 \text{ nm}$$
$$\sigma = 1.57$$

$$\frac{dN}{dD} \rightarrow N = 3140 \text{ cm}^{-3}$$
$$\mu = 164.41 \text{ nm}$$
$$\sigma = 1.28$$



Modal aerosol schemes

Twin Otter data (black)



$$\frac{dN}{dD} = \frac{dN}{dD} + \frac{dN}{dD} + \frac{dN}{dD}$$



Modal aerosol schemes



→ **MADE and MAM**

Sectional aerosol schemes

Twin Otter data (black)



→ MOSAIC, volcanic ash, (GOCART)

GOCART aerosol module

- Georgia Tech/Goddard **G**lobal **O**zone **C**hemistry **A**aerosol **R**adiation and **T**ransport model (Chin et al., JGR, 2000)
 - **Bulk aerosol:**
 - ◆ Hydrophobic black carbon (fresh soot)
 - ◆ Hydrophilic black carbon (aged/coated soot)
 - ◆ Hydrophobic organic carbon (fresh burnt biomass)
 - ◆ Hydrophilic organic carbon (aged/coated burnt biomass)
 - Fresh → aged conversion time 2.5 days
 - ◆ Other GOCART primary PM_{2.5}
 - ◆ Other GOCART primary PM₁₀
 - ◆ Sulfate (only secondary aerosol species)
 - **Sectional scheme for dust and sea salt:**
 - ◆ Dust: 0.5, 1.4, 2.4, 4.5, 8.0 μm effective radius
 - ◆ Sea salt: 0.3, 1.0, 3.2, 7.5 μm effective radius

GOCART aerosol module

GOCART comes with sulfur gas phase chemistry:

- $\text{DMS} + \text{OH} \rightarrow \text{SO}_2 + \dots$
- $\text{DMS} + \text{OH} \rightarrow \text{MSA} + \dots$
- $\text{DMS} + \text{NO}_3 \rightarrow \text{SO}_2 + \dots$
- $\text{SO}_2 + \text{OH} \rightarrow \text{SO}_4^- + \dots$

Extended gas phase chemistry can be used:

- MOZART (with KPP)
- RACM (with KPP)
- RADM (with and without KPP)

GOCART aerosol module

- **Interaction with radiation:**
 - Direct effect for some model setups
 - Effect on photochemistry
- **Interaction with clouds:**
 - Aqueous chemistry
 - ◆ $\text{SO}_2 + \text{H}_2\text{O}_2 \rightarrow \text{SO}_4^-$
 - ◆ $\text{SO}_2 + \text{O}_3 \rightarrow \text{SO}_4^-$
 - No indirect effect
 - No wet scavenging/deposition
- **No secondary organic aerosol (SOA)**

MADE aerosol module

Modal **A**erosol **D**ynamics Model for **E**urope

(Ackermann et al., Atm. Env., 1998)

- **3 log-normal aerosol modes: Aitken, accumulation, coarse**
- Mode width σ is fixed
- Aerosol number and mass variable
- (Currently no nucleation mode)
- **Interaction with radiation:**
 - Direct aerosol effect
 - Effect on photolysis
- **Interaction with clouds:**
 - Aerosol number determines cloud drop number and size
 - Radiative response → 1st indirect aerosol effect
 - ◆ only for resolved clouds (Sc)
 - Aqueous chemistry
 - Wet removal (scavenging)

MADE aerosol module

Aitken and accumulation modes:

- $\text{SO}_4^{=}$, NH_4^{+} , NO_3^{-} , H_2O
- NaCl (sea salt)
- Anthropogenic SOA from oxidation of ...
 - Alkanes
 - Alkenes
 - Aromatics
- Biogenic SOA from oxidation of ...
 - Alpha-pinene
 - Limonene
 - Isoprene
- Anthropogenic POA
- Elemental carbon (soot)
- Primary PM_{2.5}

MADE aerosol module


Coarse mode:

- **Anthropogenic primary aerosol – e.g. from**
 - Coal combustion
 - Cement manufacturing
 - Metallurgy
 - Waste incineration
- **Sea salt**
- **Soil derived particles (mineral dust)**

MADE aerosol coupling with chemistry


- **Gas phase chemistry:**
 - **RADM2** (**R**egional **A**cid **D**eposition **M**odel version **2**)
 - **RACM** (**R**egional **A**tmospheric **C**hemistry **M**echanism)
 - **RACM** NOAA/ESRL version
 - **CBMZ** (**C**arbon-**B**ond **M**echanism version **Z**)
- **Gas phase/particle partitioning (aerosol chemistry):**
 - **MARS** (**M**odel for an **A**erosol **R**eacting **S**ystem)
 - **SORGAM** (**S**econdary **O**rganic **A**erosol **M**odel)
 - **VBS** (**V**olatility **B**asis **S**et)
- **Aqueous chemistry:**
 - CMU aqueous chemistry
 - CMAQ (EPA) aqueous chemistry
 - Only for Aitken and accumulation mode
 - Only for selected gas phase chemistry options

MADE and MARS: Inorganic aerosol chemistry




MARS (**M**odel for an **A**erosol **R**eacting **S**ystem),
Saxena et al., Atm. Env., 1986


MADE/SORGAM



MADE/VBS (Volatility Basis Set)



MADE/VBS (Volatility Basis Set)



Organic aerosol mass in the surface layer
(August - September 2006)

Ahmadov et al., JGR 2012

MADE and aqueous chemistry

- **CMU aqueous chemistry** (Fahey & Pandis, Atm. Env., 2001)
 - Slow
 - Only for resolved clouds (Sc)
 - Does not really conserve mass
- **CMAQ aqueous chemistry** (Walcek & Taylor, JAS, 1986)
 - Relatively fast
 - In both resolved (Sc) and parameterized (Cu) clouds
 - Coupled to wet deposition of SO_4^- and NO_3^-

MADE + CMAQ aqueous chemistry

+ wet deposition of SO_4^- and NO_3^- :

- **chem_opt = 41 : RADM2 gas phase chemistry**


Planned for WRF/Chem 3.5.1:

- **chem_opt = 42 : RACM gas phase chemistry (KPP)**
- **chem_opt = 43 : RACM ESRL gas phase chemistry (KPP)**

MADE and CMAQ aqueous chemistry


Details of aqueous chemistry depend on cloud type

Stratocumulus




~ 10km

Cumulus




~100m

Cumulonimbus



~1km

MADE and wet deposition



Precipitation	
r	model/obs.
0.81	1.25

May-September 2006
(National Atmospheric Deposition Program)

SO ₄ ⁼ wet dep.	
r	model/obs.
0.86	0.53

MOSAIC aerosol module

Model for Simulating Aerosol Interactions and Chemistry
(Zaveri et al., JGR, 2008)

- Modern aerosol scheme in WRF/Chem
- 4 or 8 aerosol size sections (bins) 39 nm – 10 μm
- (Lower bin boundary of 39 nm too large for nucleation)
- Interaction with radiation:
 - Direct aerosol effect
 - Effect on photolysis
- Interaction with clouds:
 - Aerosol number determines cloud drop number and size
 - Radiative response \rightarrow 1st indirect aerosol effect
 - Aqueous chemistry
 - Wet removal (scavenging)
 - only for resolved clouds (Sc)

MOSAIC aerosol module


Aerosol composition

- $\text{SO}_4^{=}$, NH_4^{+} , NO_3^{-} , H_2O
- NaCl (sea salt)
- CH_3SO_3 (methanesulfonate)
- carbonate (CO_3)
- calcium (Ca)
- black carbon (BC)
- primary organic mass (OC)
- other inorganic mass (minerals, trace metals)

MOSAIC aerosol coupling with chemistry

- **Gas phase chemistry:**
 - **CBMZ** (**C**arbon-**B**ond **M**echanism version **Z**)
 - ◆ “Standard” gas phase chemical scheme for MOSAIC
 - **SAPRC99** (extensive VOC chemistry)
 - ◆ Works with the VBS SOA scheme
 - **MOZART** (**M**odel for **O**zone and **R**elated chem. **T**racers)
 - ◆ Works with the VBS SOA scheme
- **Gas phase/particle partitioning (aerosol chemistry):**
 - **MTEM** (**M**ulticomponent **T**aylor **E**xpansion **M**ethod)
 - **MESA** (**M**ulticomponent **E**quilibrium **S**olver for **A**erosols)
 - **VBS** (**V**olatility **B**asis **S**et)
- **Aqueous chemistry:**
 - CMU aqueous chemistry, only for resolved clouds (Sc)
 - Not with KPP versions of gas phase chemistry schemes

MOSAIC, MTEM, and MESA



MTEM calculates activity coefficients

MESA solves ion-equilibria in the liquid phase

For SOA: VBS (Volatility Basis Set) scheme

MTEM (Multicomponent Taylor Expansion Method), Zaveri et al., JGR 2005a

MESA (Multicomponent Equilibrium Solver for Aerosols), Zaveri et al., JGR 2005b

MAM aerosol module

Modal Aerosol Model from CAM5

- New in WRF/Chem 3.5
- **3 or 7 log-normal aerosol modes: MAM3 and MAM7**
- Mode width σ is fixed
- Aerosol number and mass variable
- Liu et al., Geosci. Model Dev., 5, 709-739, 2012

MAM 3


Aitken mode



- Sulfate ($\text{SO}_4^{=}$)
- SOA
- Sea salt

15 – 53 nm


Accumulation mode



- Sulfate ($\text{SO}_4^{=}$)
- SOA
- Primary organic matter
- Sea salt
- Black carbon
- Mineral dust
- Sea salt

58 – 270 nm

Coarse mode



- Sulfate ($\text{SO}_4^{=}$)
- Mineral dust
- Sea salt

0.8 – 3.65 μm


Dry particle diameter

MAM 7


Aitken mode

Accumulation mode

Coagulation,
condensation


- 
- Sulfate (SO_4^-)
 - SOA
 - Sea salt

Primary carbon


- 
- Sulfate (SO_4^-)
 - SOA
 - Primary organic matter
 - Sea salt
 - Black carbon

- 
- Primary organic matter
 - Black carbon


Fine sea salt, SO_4^-




Coarse sea salt, SO_4^-



Fine dust, SO_4^-



Coarse dust, SO_4^-



MAM aerosol module

- **Currently only one gas phase chemistry scheme**
 - **CBMZ** (**C**arbon-**B**ond **M**echanism version **Z**)
- **Interaction with radiation:**
 - Coupled to RRTMG radiation → Direct aerosol effect

As in CAM5:

- **Gas phase/particle partitioning (aerosol chemistry):**
 - Condensation of water vapor and of the 4 inorganic trace gas species: NH_3 , H_2SO_4 , HNO_3 , HCl
- **Interaction with clouds only resolved clouds (Sc):**
 - Coupled to Morrison & Gettelman cloud microphysics
 - Radiative response → 1st indirect aerosol effect
 - Wet removal (scavenging)
 - Aqueous chemistry
- **Dry deposition**

Volcanic ash

- **10 bins for volcanic ash aerosol**
- **Transport, settling, dry deposition**
- **Currently no other aerosol**
- **Single active volcano**
- **Database of 1535 volcanoes (latitude, longitude, height)**

How to tell WRF/Chem what to do

../WRFV3/test/em_real/real.exe

| ../WRFV3/test/em_real/namelist.input

../WRFV3/test/em_real/...

../WRFV3/test/em_real/...

...

...

&chem

chem_opt = **42**

photdt = 0.25

chemdt = 0

...

aerchem_onoff = **1**

...

conv_tr_aqchem = **1**

MADE/SORGAM,
RACM, CMAQ
aqueous chemistry

Switches all aerosol
processes on/off

CMAQ aqueous
chemistry on in Cu

Resources

- **WRF/Chem User's Guide**
 - Model options (namelist parameters)
 - Combinations of physical/chemical schemes
 - ...
- **Papers referenced in the WRF/Chem User's Guide**
- **WRF/Chem source code**
- **WRF/Chem Help (wrfchemhelp.gsd@noaa.gov)**
- **Yours truly (jan.kazil@noaa.gov)**