

Emissions Data Tools for Use with WRF-Chem

Steven Peckham

Emissions Data

- Several tools are available to help generate emissions
 - **Prep_chem_sources**
 - Anthropogenic
 - Biogenic (included in anthropogenic)
 - Biomass Burning
 - GOCART Background fields
 - Volcano
 - Anthro_emiss (NCAR/ACD) – limited testing (MOZART & CAM)
 - **MEGAN** (NCAR/ACD)
 - Biogenic emissions
 - FINN (NCAR/ACD)
 - Biomass Burning (retrospective simulation)
 - Emiss_v03 (NOAA)
 - Anthropogenic (USA only)
 - Others
 - SMOKE (US EPA)

Generating Emissions Fields for WRF-Chem with PREP-CHEM-SRC

Rafael Santos Lima, Marcelo Alonso, Megan Bela,
Valter Oliveira, Rafael Fonseca, Madeleine Gácita, Gabriel Pereira,
Karla M. Longo, Saulo R. Freitas, Georg Grell, Steven Peckham

megan.bela@colorado.edu
gmai@cptec.inpe.br
meioambiente.cptec.inpe.br



PREP-CHEM-SRC

Emissions Utility for:

Anthropogenic

Biogenic

Biomass burning and plume rise

Volcanoes

How to generate emissions

Compiling

Namelist

Running PREP-CHEM-SRC and `convert_emiss`

Anthropogenic emissions

Global Inventories

RETRO ($0.5^\circ \times 0.5^\circ$, monthly, 1960-2000)

EDGAR v4.2 ($0.1^\circ \times 0.1^\circ$, annual, 1970-2008)


CO_2 , CH_4 , N_2O , HFCs, PFCs, SF_6

GOCART

OC, BC and SO_2 ($1^\circ \times 1^\circ$, annual, 2006)


DMS ($1^\circ \times 1.25^\circ$, monthly)

NO_3 , H_2O_2 and OH (3D, $1^\circ \times 1.25^\circ$
monthly, 2006)





Anthropogenic emissions

Example for Alaska




Anthropogenic emissions

AREA DELIMITER algorithm distributes emissions on high resolution grids



Anthropogenic emissions

South America: Updated local inventories and extrapolation to cities without inventories based on socioeconomic data



CO emissions ($\times 10^6$ kg m^2 day $^{-1}$) on a 20 km grid covering South America without (A) and with (B) updated inventories

Biogenic emissions (if bio_emiss_opt=0)

1) GEIA


1° x 1°, monthly, 2002

Acetone, C₂H₄, C₂H₆, C₃H₆,
C₃H₈, CO, CH₃OH, DMS, NO,
Isoprene, Terpenes and NVOC

2) MEGAN 2000 climatology

0.5° x 0.5°, monthly, 2000

CO, CH₄, C₂H₄, C₂H₆, C₃H₆, C₃H₈,
CH₃OH, Formaldehyde, Acetaldehyde,
Acetone, other Ketones, Toluene,
Isoprene, Monoterpenes and
Sesquiterpenes



Daily emissions from (A) GEIA (B) 3BEM (C) RETRO for 27 August 2002 on a 0.2° grid


Alonso et al. (2010)

Biomass burning in the developing world


Emissions_Yevich_Logan

$10^0 \times 10^0$, Tg dry matter yr⁻¹


Woodfuel (fuelwood and charcoal) use



Crop residue and dung use




Burning of agricultural residue in the fields



Biomass burning emissions

Brazilian Biomass Burning
Emission Model (**3BEM**)
Model resolution, daily

Global Fire Emissions Database (**GFEDv2**)
 $1^\circ \times 1^\circ$, 8-day or monthly, 1997 - 2004




Average daily CO emissions, Aug.-Oct. 2002, 35 km

Freitas et al. (2011)

Wildfire Satellite Data


- Moderate-Resolution Imaging Spectroradiometer (MODIS) instruments on NASA's Aqua and Terra satellites.
 - Daily global 1km fire data
 - Available in real-time
 - <https://earthdata.nasa.gov/data/near-real-time-data/firms>
 - Data archives can be found online
 - Reads text (.txt) data files
- GOES, MET, MTSAT
 - Automated Biomass Burning Algorithm (ABBA) fire products (<http://wfabba.ssec.wisc.edu>)
 - Derived from radiances from bands 1 (visible), 2 (3.9 micron), and 4 (11 micron)
 - Available in real time (GOES: North & South America domain, MET-9: Africa, MTSAT: Australia)




3BEM

Biomass burning emissions inventory Regional scale – daily basis

density of carbon data




near real time fire product



6 types of biomes
110 chemical species

land use data



Andreae and Merlet, 2001
emission & combustion factors

Biome category	Emission Factor for CO (g/kg)	Emission Factor for PM2.5 (g/kg)	Aboveground biomass density (α, kg/m ²)	Combustion factor (β, fraction)
Tropical forest ¹	110.	8.3	20.7	0.48
South America savanna ²	63.	4.4	0.9	0.78
Pasture ³	49.	2.1	0.7	1.00

¹ Average values for primary and second-growth tropical forests, ² Average values for campo cerrado (C3) and cerrado sensu stricto (C4), ³ value for campo limpo (C1). All numbers are from Ward et al.,

mass estimation


$$M_{[\eta]} = \alpha_{veg} \cdot \beta_{veg} \cdot E_{f_{veg}}^{[\eta]} \cdot a_{fire}$$

CO source emission (kg m⁻²day⁻¹)

3BEM Plume Rise

Biomass burning
and wildfires


Smoldering : mostly surface emission.
Flaming: mostly direct injection in the PBL,
free troposphere or stratosphere.



Plume rise model

total emission flux: F_η being λ the smoldering fraction



$$\text{smoldering term : } E_\eta = \frac{\lambda F_\eta}{\rho_{air} \Delta z_{\text{first phys. model layer}}}$$

$$\text{flaming term : } E_\eta = \frac{(1 - \lambda) F_\eta}{\rho_{air} \Delta z_{\text{injection layer}}}$$


Example in
the model:

flaming
emission

smoldering
emission



diurnal cycle of the
burning for S. America:

$$E_\eta(t) = r(t) E_\eta$$

Freitas et al. (2011)


Recent Improvements: Environmental Wind Effects on Plume Rise



Biomass burning plumes in the Amazon region
without (left) and with (right) environmental wind shear

Photos: M.O. Andreae, M. Welling

Environmental Wind Effects on Plume Rise




$$\lambda_{\text{entr}} = \frac{2\alpha}{R} |w|$$

$$\delta_{\text{entr}} = \frac{2}{\pi R} (u_e - u)$$

W : vertical velocity
 VMD : vertical mass distribution
 Ea : Entrainment acceleration
 Ba : buoyancy acceleration
 CW : total condensate water


1-D PRM results for a 50 ha fire, calm and windy conditions



Freitas et al. (2010)

Volcano emissions

Based on Mastin et al. (2009) database of 1535 volcanoes
Mass eruption rate, plume height and time duration
SO₂ from AEROCOM program, 1979 – 2007 (Diehl, 2009)



SO₂ emissions on 27 August 2002 on a 0.2° rectangular projection
grid: (A) Diehl (2009), (B) 3BEM, (C) EDGAR

Freitas et al. (2011)

Compiling PREP-SRC-CHEM

- Required libraries: HDF4/HDF5, zlib, jpeg, netCDF
- cd to:
`PREP-CHEM-SRC-1.4/bin/build`
- Set library paths in:
`include.mk.<compiler>`
- Compile
`make OPT=<compiler> CHEM=RADM_WRF_FIM`
- cd to:
`PREP-CHEM-SRC-1.4/bin`

Executable : *prep_chem_sources_RADM_WRF_FIM.exe*

Input file (namelist): *prep_chem_sources.inp*

Input file (namelist): prep_chem_sources.inp

\$RP_INPUT

!----- **grid_type**

grid_type= 'lambert', ! 'polar' = polar stereo. grid output
 ! 'll' = lat/lon grid output
 ! 'lambert' = lambert grid output
 ! 'mercator' = mercator grid output

!----- **date of emission**

ihour=0, iday=12, imon=7, iyear=2004,

!----- **select the sources datasets to be used: 1 = yes, 0 = not**

use_retro=1,
retro_data_dir='/DATA/Emission_data/RETRO/anthro',
use_edgar =2, ! 0 - not, 1 - Version 3, 2 - Version 4 for some species
use_seac4rs=1,
ue_streets = 0,

use_gocart=1,
user_data_dir='/DATA/EMISSION_DATA/SouthAmerica_Megacities',

use_bioge =2, ! 1 - GEIA, 2 - MEGAN
use_fwbawb=1,
fwbawb_data_dir='/DATA/Emission_data/Emissions_Yevich_Logan',
use_gfedv2=0,
use_bbem=1,
use_bbem_plumerise=1,

Input file (namelist): prep_chem_sources.inp

!----- **if the merging of gfedv2 with bbem is desired (=1, yes, 0 = no)**

merge_GFEDv2_bbem =0,

!----- **Fire product for 3BEM/3BEM-plumerise emission models**

bbem_wfabba_data_dir='/DATA/Emission_data/fires_data/WF_ABBA/filt/f,

bbem_modis_data_dir ='/DATA/Emission_data/fires_data/MODIS/Fires.',

bbem_inpe_data_dir ='/DATA/Emission_data/fires_data/DSA/Focos',

bbem_extra_data_dir ='/DATA/Emission_data/fires_data/xxxxx,

!----- **gocart background**

use_gocart_bg=1,

!----- **volcanoes emissions**

use_volcanoes=0,

volcano_index=0, !REDOUBT

use_these_values='NONE',

! define a text file for using external values for INJ_HEIGHT, DURATION,

! MASS ASH (units are meters - seconds - kilograms) and the format for

! a file 'values.txt' is like this: 11000. 10800. 1.5e10

! use_these_values='values.txt',

begin_eruption='198912141930', !begin time UTC of eruption YYYYMMDDhhmm

!----- degassing volcanoes emissions

use_degass_volcanoes=0,

degass_volc_data_dir='/DATA/EMISSION_DATA/VOLC_SO2',

Input file (namelist): prep_chem_sources.inp



!----- **For regional grids (polar, Lambert, Mercator)**

```
NGRIDS = 3,          ! Number of grids to run
NNXP   = 391,463,499, ! Number of x gridpoints
NNYP   = 271,454,478, ! Number of y gridpoints
NXTNEST = 0, 1, 2,    ! Grid number which is the next coarser grid
DELTAX  = 18000,
DELTAY  = 18000,      ! X and Y grid spacing
! Nest ratios between this grid and the next coarser grid.
NSTRATX = 1, 3, 3,    ! x-direction
NSTRATY = 1, 3, 3,    ! y-direction
NINEST  = 1, 78, 128, ! Grid point on the next coarser
NJNEST  = 1, 30, 153, ! nest where the lower southwest
! NKNEST = 1, 1, 1,   ! nest where the lower southwest
                ! corner of this nest will start.
                ! If NINEST or NJNEST = 0, use CENTLAT/LON
POLELAT = 15.,        ! If polar, latitude/longitude of pole point
POLELON = 10.,        ! If lambert, lat/lon of grid origin (x=y=0.)
STDLAT1 = 0.,         ! If polar, unused
STDLAT2 = 15.,        ! If lambert, standard latitudes of projection (truelat2/truelat1 from
                    namelist.wps, STDLAT1 < STDLAT2)
CENTLAT  = 15.0,
CENTLON  = 10.0,
```

Running PREP-CHEM-SRC and convert_emiss


prep_chem_sources_RADM_WRF_FIM.exe

./real.exe
(chem_opt=0,)




Binary emissions (**-ab.bin*,
**-bb.bin*, **gocartBG.bin*, **volc.bin*)

wrfinput_d01



../chem/convert_emiss.exe



netCDF emissions (*wrfchemi**,
*wrffirechemi**, *wrfchemi_gocart_bg_**)

References

Alonso, M. F. ; Longo, K. M. ; Freitas, S. R. ; Fonseca, R. M. ; Marecal, V. ; Pirre, M. ; Gallardo, L. . An urban emissions inventory for South America and its application in numerical modeling of atmospheric chemical composition at local and regional scales. *Atmospheric Environment*, v. 44, p. 5072-5083, 2010.

Freitas, S. R. ; Longo, K. M. ; Alonso, M. F. ; Pirre, M. ; Marecal, V. ; Grell, G. ; Stockler, R. ; Mello, R. F. ; Sánchez Gácita, M. . PREP-CHEM-SRC 1.0: a preprocessor of trace gas and aerosol emission fields for regional and global atmospheric chemistry models. *Geoscientific Model Development*, v. 4, p. 419-433, 2011.

Freitas, S. R. , Longo, Karla , Trentmann, J. , Latham, D. Technical Note: Sensitivity of 1-D smoke plume rise models to the inclusion of environmental wind drag. *Atmospheric Chemistry and Physics*, v. 10, p. 585-594, 2010.

Freitas, S. R., K. M. Longo, R. Chatfield, D. Latham, M. A. F. Silva Dias, M. O. Andreae, E. Prins, J. C. Santos, R. Gielow and J. A. Carvalho Jr.: Including the sub-grid scale plume rise of vegetation fires in low resolution atmospheric transport models. *Atmospheric Chemistry and Physics*, v. 7, p. 3385-3398, 2007.

Freitas, S. R.; Longo, K. M.; M. Andreae. The impact of including the plume rise of vegetation fires in numerical simulations of associated atmospheric pollutants. *Geophys. Res. Lett.*, 33, L17808, doi:10.1029/2006GL026608, 2006.

Yevich, R. and J.A. Logan, An assessment of biofuel use and burning of agricultural waste in the developing world, *Global Biogeochemical Cycles*, 2003




Questions?

wrfchemhelp.gsd@noaa.gov
gmai@cptec.inpe.br
megan.bela@colorado.edu

Using MEGAN with WRF-Chem

Gabriele Pfister, Stacy Walters, Christine
Wiedinmyer, Alex Guenther, Mary Barth,
Louisa Emmons, Tiffany Duhl

Preprocessors available from: www.acd.ucar.edu



WRF-Chem

WRF-Chem is the Weather Research and Forecasting (WRF) model coupled with Chemistry. The model simulates the emission, transport, mixing, and chemical transformation of trace gases and aerosols simultaneously with the meteorology. The model is used for investigation of regional-scale air quality, field program analysis, and cloud-scale interactions between clouds and chemistry.

The development of WRF-Chem is a collaborative effort among the community. NOAA/ESRL scientists are the leaders and caretakers of the code. The [Official WRF-Chem web page](#) is located at the NOAA web site. Our model development is closely linked with both NOAA/ESRL and DOE/PNNL efforts. Description of [PNNL WRF-Chem model](#) development is located at the PNNL web site as well as the [PNNL Aerosol Modeling Testbed](#).

Use the [MOZART Download](#) page to retrieve MOZART-4 model results.

Information on running WRF-Chem with the MOZART chemical mechanism can be found in the [MOZCART User's Guide](#).

Processors Available to the Community:

NEW (November 2011): Preprocessors have been updated to work for lat/lon projections in addition to Lambert, Mercator and Polar. The mozbc tool has been updated to enable time interpolation.

mozbc

NCAR/ACD has developed a program to create time-varying chemical boundary conditions for WRF-Chem from MOZART-4 output. For questions about running mozbc please contact: Stacy Walters (stacy@ucar.edu), Mary Barth (barthm@ucar.edu), or Gabriele Pfister (pfister@ucar.edu). To obtain mozbc, see the **Download** section below.

bio_emiss

Bio_emiss is a pre-processor for creating MEGAN input for WRF-Chem. To obtain bio_emiss, see the **Download** section below.

preprocessor tools

Pre-processor tools for running WRF-Chem / MOZCART. See the **Download** section below.


Download

- Preprocessors are in FORTRAN
- README details compilation and execution
- Require netcdf libraries; work on Portland Group or IBM fortran 90 compilers
- Domain information derived from `wrfinput_d<domain>`

MEGAN online biogenic emissions

Model of Emissions of Gases and Aerosols from Nature


- Estimate emissions of VOCs, NO_x and CO from vegetation
- Driving variables include landcover, LAI, weather, and atmospheric chemical composition
- Reference: *Guenther et al.*, GMD 2012 (for MEGAN v2.1)
- Note: as of current land cover used in MEGAN differs from that used within WRF-Chem
- Planned:
 - Update to latest MEGAN current version in WRF-Chem: 2.04
 - link to WRF-Chem land cover/CLM



from Guenther et al., 2012

MEGAN preprocessor

- Static input fields needed to run with online MEGAN biogenic emissions: *Isoprene Emissions Factors, past/current conditions (monthly LAI, solar radiation & temperature), Fractional coverage of broadleaf and needelleaf trees, shrubs and herbaceous*
- Compatible with MOZART, CBMZ, RADM, RACM, SAPRC (see `module_data_mgn2mech.F` for species mapping)
- Download source code (`megan_bio_emiss.tar`) and global input data (`megan.data.tar.gz`)
- `megan_bio_emiss` is a **single cpu code**, which:
 - ✓ reads global MEGAN input data
 - ✓ maps them on the WRF-Chem domain
 - ✓ creates `wrfbiochemi_d<domain>` file



bio_emiss

Bio_emiss is a pre-processor for creating MEGAN input for WRF-Chem. To obtain bio_emiss, see the **Download** section below.

MEGAN preprocessor

- To compile:
make_util megan_bio_emiss - creates the executable *megan_bio_emiss*
- *megan_bio_emiss* is controlled by a namelist file
e.g. “*megan_bio_emiss.inp*”

&control

```
domains = 3,  
start_lai_mnth = 4,  
end_lai_mnth = 6,  
wrf_dir = '/home/me/megan/wrf_files',  
megan_dir = '/home/me/megan/30sec'  
/
```

*creates wrfbiochemi_dnn for three domains (default: 1)
starting month for the monthly LAI (default: 1)
ending month for the monthly LAI (default: 12)
path to wrfinput_dnn (default: current)
path to MEGAN input files (default: current)*

- To run : *megan_bio_emiss < megan_bio_emiss.inp > megan_bio_emiss.out*

Running WRF-Chem with MEGAN

- WRF-Chem output variables: EBIO_<species>, ...
- namelist.input:

```
&time_control
```

```
(activate settings only either during real.exe or initial wrf.exe)
```

```
auxinput6_interval_h      = 24
```

```
auxinput6_inname         = 'wrfbiochemi_d01',
```


```
io_form_auxinput6        = 2,
```

```
&chem
```

```
bio_emiss_opt = 3
```

```
bioemdt = your choice (minutes)
```

```
ne_area = # chemical species
```



EBIO_ISO, 20 July 2008 21 UTC

For Questions:

Gabriele Pfister *pfister@ucar.edu*

Stacy Walters *walters@ucar.edu*

