



Chemical mechanisms and Kinetic PreProcessor (KPP) in WRF-Chem 3.7

Ravan Ahmadov

ravan.ahmadov@noaa.gov


Cooperative Institute for Research in Environmental Sciences, CU Boulder
Earth System Research Laboratory, NOAA, Boulder

Acknowledgements: S.McKeen, G.Grell, S.Peckham, S.-W.Kim, M.Salzmann

**WRF-Chem tutorial,
Boulder, August 3, 2015**

Outline

- ❑ **Chemical mechanisms in the WRF-Chem V3.7 model**
- ❑ **Kinetic PreProcessor (KPP) and WRF-Chem KPP Coupler (WKC)**
- ❑ **Adding chemical mechanisms to the WRF-Chem model using KPP**
- ❑ **Additional notes**



The gas phase chemistry mechanisms in WRF-Chem3.6

- ❑ Regional Acid Deposition Model, 2nd generation (**RADM2**)
- ❑ Regional Atmospheric Chemistry Mechanism (**RACM**)
- ❑ RACM - Mainz Isoprene Mechanism (**RACM-MIM**)
- ❑ RACM - Earth System Research Laboratory (**RACM-ESRL**), *updated RACM-MIM*
- ❑ Carbon Bond Mechanism (**CB4**)
- ❑ Carbon-Bond Mechanism version Z (**CBMZ**)
- ❑ Model of Ozone and Related Chemical Tracers (**MOZART**)
- ❑ Nonmethane Hydrocarbons Chemistry (**NMHC9**)
- ❑ Statewide Air Pollution Research Center (**SAPRC99**)
- ❑ **CRIMECH**

different implementations, coupled to different aerosol schemes and aqueous chemistry, suitable for different applications ranging from regional air quality to global atmospheric chemistry simulations (no halogen chemistry yet!)

How complex are the gas chemistry options in WRF-Chem?

- **Passive tracer schemes (chem_opt=13-15): no chemistry, transport only, anthropogenic emissions only**
- **Passive tracer schemes for greenhouse gases (chem_opt=16, 17): no chemistry, transport only, anthropogenic emissions and biospheric fluxes of CO₂ and CH₄**
- **Computationally efficient options using the RADM scheme (e.g. chem_opt=1, 2, 101): for air quality modeling, not all biogenic VOCs are included**
- **Options based on the RACM schemes (e.g. chem_opt=102-105): more updated version of RADM, more biogenic VOCs are included, updated isoprene chemistry, reaction rates**

How complex are these gas chemistry options?

- **Passive tracer schemes (chem_opt=13-15): no chemistry, transport only, anthropogenic emissions only**
- **Passive tracer schemes for greenhouse gases (chem_opt=16, 17): no chemistry, transport only, anthropogenic emissions and biospheric fluxes of CO₂ and CH₄**
- **Computationally efficient options using the RADM scheme (e.g. chem_opt=1, 2, 101): for air quality modeling, not all biogenic VOCs are included**
- **Options based on the CRIMECH schemes (e.g. chem_opt=600, 601): more updated version of RADM, more biogenic VOCs are included, updated isoprene chemistry, reaction rates**

Some references

- ❑ Stockwell, W. R., et al. (1997), A new mechanism for regional atmospheric chemistry modeling, *J. Geophys. Res.-Atmos.*, 102(D22), 25847-25879.
- ❑ Stockwell, W. R., et al. (1990), The 2nd generation regional acid deposition model chemical mechanism for regional air-quality modeling, *J. Geophys. Res.-Atmos.*, 95(D10), 16343-16367.
- ❑ Horowitz, L. W., et al. (2003), A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *J. Geophys. Res.*, 108(D24), 4784, doi:4710.1029/2002JD002853.
- ❑ Gross, A., and W. R. Stockwell (2003), Comparison of the EMEP, RADM2 and RACM mechanisms, *Journal of Atmospheric Chemistry*, 44(2), 151-170.
- ❑ Geiger, H., et al. (2003), The tropospheric degradation of isoprene: an updated module for the regional atmospheric chemistry mechanism, *Atmos. Environ.*, 37(11), 1503-1519.
- ❑ Luecken, D. J., et al. (2008), Effects of using the CB05 vs. SAPRC99 vs. CB4 chemical mechanism on model predictions: Ozone and gas-phase photochemical precursor concentrations, *Atmos. Environ.*, 42(23), 5805-5820.
- ❑ Cai et al. (2011), Photochemical Modeling in California with Two Chemical Mechanisms: Model Intercomparison and Response to Emission Reductions, *J. Air & Waste Manage. Assoc.*
- ❑ Peckham S. et al. (2013), WRF-CHEM 3.5 User's Guide.

Chemistry mechanisms in WRF-Chem3.6

<i>Chemical mechanisms</i>	<i>Fixed versions</i>	<i>KPP</i>	<i>Coupled to the aerosol schemes</i>
RADM2	Yes	Yes	MADE/SORGAM, GOCART
RACM	None	Yes	MADE/SORGAM, GOCART
RACM-MIM	None	Yes	None
RACM-ESRL	None	Yes	MADE/SORGAM, MADE/ SOA_VBS
CB4	None	Yes	None
CBMZ	Yes	Yes	MOSAIC
MOZART	None	Yes	GOCART
SAPRC99	None	Yes	MOSAIC
NMHC9	None	Yes	None
CRIMECH	None	Yes	MOSAIC

Table 1. RACM Mechanism Species List

No.	Species	Definition	Carbon Number	Molecular Weight
<i>Stable Inorganic Compounds</i>				
Oxidants				
1	O ₃	ozone		48
2	H ₂ O ₂	hydrogen peroxide		34
Nitrogenous compounds				
3	NO	nitric oxide		30
4	NO ₂	nitrogen dioxide		46
5	NO ₃	nitrogen trioxide		62
6	N ₂ O ₅	dinitrogen pentoxide		108
7	HONO	nitrous acid		47
8	HNO ₃	nitric acid		63
9	HNO ₄	pernitric acid		79
Sulfur compounds				
10	SO ₂	sulfur dioxide		64
11	SULF	sulfuric acid		98
Carbon oxides				
12	CO	carbon monoxide	1	28
13	CO ₂	carbon dioxide	1	44
<i>Abundant Stable Species</i>				
14	N ₂	nitrogen		28
15	O ₂	oxygen		32
16	H ₂ O	water		18
17	H ₂	hydrogen		2
<i>Inorganic Short-Lived Intermediates</i>				
18	O ³ P	ground state oxygen atom, O(³ P)		16
19	O ¹ D	excited state oxygen atom, O(¹ D)		16
Odd hydrogen				
20	HO	hydroxy radical		17
21	HO ₂	hydroperoxy radical		33

Table 2b. The RACM Mechanism

Reaction No.	Reaction	A , $\text{cm}^3 \text{s}^{-1}$	E/R , K	k^a
<i>Inorganic Reactions</i>				
(R24)	$\text{O}^3\text{P} + \text{O}_2 \rightarrow \text{O}_3$	Table 2f		1.50×10^{-14}
(R25)	$\text{O}^3\text{P} + \text{O}_3 \rightarrow 2 \text{O}_2$	8.00×10^{-12}	2060	7.96×10^{-15}
(R26)	$\text{O}^1\text{D} + \text{N}_2 \rightarrow \text{O}^3\text{P} + \text{N}_2$	1.80×10^{-11}	-110	2.60×10^{-11}
(R27)	$\text{O}^1\text{D} + \text{O}_2 \rightarrow \text{O}^3\text{P} + \text{O}_2$	3.20×10^{-11}	-70	4.05×10^{-11}
(R28)	$\text{O}^1\text{D} + \text{H}_2\text{O} \rightarrow \text{HO} + \text{HO}$	2.20×10^{-10}		2.20×10^{-10}
(R29)	$\text{O}_3 + \text{HO} \rightarrow \text{HO}_2 + \text{O}_2$	1.60×10^{-12}	940	6.83×10^{-14}
(R30)	$\text{O}_3 + \text{HO}_2 \rightarrow \text{HO} + 2 \text{O}_2$	1.10×10^{-14}	500	2.05×10^{-15}
(R31)	$\text{HO} + \text{HO}_2 \rightarrow \text{H}_2\text{O} + \text{O}_2$	4.80×10^{-11}	-250	1.11×10^{-10}
(R32)	$\text{H}_2\text{O}_2 + \text{HO} \rightarrow \text{HO}_2 + \text{H}_2\text{O}$	2.90×10^{-12}	160	1.70×10^{-12}
(R33)	$\text{HO}_2 + \text{HO}_2 \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$	Table 2f		2.92×10^{-12}
(R34)	$\text{HO}_2 + \text{HO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + \text{O}_2 + \text{H}_2\text{O}$	Table 2f		6.58×10^{-30}
(R35)	$\text{O}^3\text{P} + \text{NO} \rightarrow \text{NO}_2$	Table 2d		1.66×10^{-12}
(R36)	$\text{O}^3\text{P} + \text{NO}_2 \rightarrow \text{NO} + \text{O}_2$	6.50×10^{-12}	-120	9.72×10^{-12}
(R37)	$\text{O}^3\text{P} + \text{NO}_2 \rightarrow \text{NO}_3$	Table 2d		1.58×10^{-12}
(R38)	$\text{HO} + \text{NO} \rightarrow \text{HONO}$	Table 2d		4.87×10^{-12}
(R39)	$\text{HO} + \text{NO}_2 \rightarrow \text{HNO}_3$	Table 2d		1.15×10^{-11}
(R40)	$\text{HO} + \text{NO}_3 \rightarrow \text{NO}_2 + \text{HO}_2$	2.20×10^{-11}		2.20×10^{-11}
(R41)	$\text{HO}_2 + \text{NO} \rightarrow \text{NO}_2 + \text{HO}$	3.70×10^{-12}	-250	8.56×10^{-12}
(R42)	$\text{HO}_2 + \text{NO}_2 \rightarrow \text{HNO}_4$	Table 2d		1.39×10^{-12}
(R43)	$\text{HNO}_4 \rightarrow \text{HO}_2 + \text{NO}_2$	Table 2e		8.62×10^{-2}
(R44)	$\text{HO}_2 + \text{NO}_3 \rightarrow 0.3 \text{HNO}_3 + 0.7 \text{NO}_2 + 0.7 \text{HO} + \text{O}_2$	3.50×10^{-12}		3.50×10^{-12}
(R45)	$\text{HO} + \text{HONO} \rightarrow \text{NO}_2 + \text{H}_2\text{O}$	1.80×10^{-11}	390	4.86×10^{-12}
(R46)	$\text{HO} + \text{HNO}_3 \rightarrow \text{NO}_3 + \text{H}_2\text{O}$	Table 2f		1.47×10^{-13}
(R47)	$\text{HO} + \text{HNO}_4 \rightarrow \text{NO}_2 + \text{O}_2 + \text{H}_2\text{O}$	1.30×10^{-12}	-380	4.65×10^{-12}
(R48)	$\text{O}_3 + \text{NO} \rightarrow \text{NO}_2 + \text{O}_2$	2.00×10^{-12}	1400	1.82×10^{-14}
(R49)	$\text{O}_3 + \text{NO}_2 \rightarrow \text{NO}_3 + \text{O}_2$	1.20×10^{-13}	2450	3.23×10^{-17}
(R50)	$\text{NO} + \text{NO} + \text{O}_2 \rightarrow \text{NO}_2 + \text{NO}_2$	3.30×10^{-39}	-530	1.95×10^{-38}
(R51)	$\text{NO}_3 + \text{NO} \rightarrow \text{NO}_2 + \text{NO}_2$	1.50×10^{-11}	-170	2.65×10^{-11}
(R52)	$\text{NO}_3 + \text{NO}_2 \rightarrow \text{NO} + \text{NO}_2 + \text{O}_2$	4.50×10^{-14}	1260	6.56×10^{-16}
(R53)	$\text{NO}_3 + \text{NO}_2 \rightarrow \text{N}_2\text{O}_5$	Table 2d		1.27×10^{-12}
(R54)	$\text{N}_2\text{O}_5 \rightarrow \text{NO}_2 + \text{NO}_3$	Table 2e		4.36×10^{-2}
(R55)	$\text{NO}_3 + \text{NO}_3 \rightarrow \text{NO}_2 + \text{NO}_2 + \text{O}_2$	8.50×10^{-13}	2450	2.29×10^{-16}
(R56)	$\text{HO} + \text{H}_2 \rightarrow \text{H}_2\text{O} + \text{HO}_2$	5.50×10^{-12}	2000	6.69×10^{-15}
(R57)	$\text{HO} + \text{SO}_2 \rightarrow \text{SULF} + \text{HO}_2$	Table 2d		8.89×10^{-13}
(R58)	$\text{CO} + \text{HO} \rightarrow \text{HO}_2 + \text{CO}_2$	Table 2f		2.40×10^{-13}

Chemistry options in WRF-Chem3.6

chem_opt =0 no chemistry

= 1 RADM2 chemical mechanism - no aerosols

= 2 RADM2 chemical mechanism and MADE/SORGAM aerosols

No indirect effect To have radiative feed back with the chemistry/aerosols use `ra_sw_physics = 2` (Goddard shortwave scheme). For dust and sea salt use `dust_opt=2, seas_opt=2`

= 5 CBMZ chemical mechanism with Dimethylsulfide

= 6 CBMZ chemical mechanism without DMS

= 7 CBMZ chemical mechanism (chem_opt=6) and MOSAIC using 4 sectional aerosol bins

No indirect effect To have radiative feed back with the chemistry/aerosols use , `ra_sw_physics = 2`, for dust and seasalt use `dust_opt=2, seas_opt=2`

= 8 CBMZ chemical mechanism (chem_opt=6) and MOSAIC using 8 sectional aerosol bins.

No indirect effect To have radiative feed back with the chemistry/aerosols use, `ra_sw_physics = 2`, for dust and seasalt use `dust_opt=2, seas_opt=2`.

= 9 CBMZ chemical mechanism (chem_opt=6) and MOSAIC using 4 sectional aerosol bins) including some aqueous reactions

For direct and indirect effect use: `phot_opt=2; ra_sw_physics=2; progn=1; mp_physics=2; aer_ra_feedback=1; wetscav_onoff=1; cldchem_onoff=1,`

For dust and sea salt use `dust_opt=2, seas_opt=2`

= 10 CBMZ chemical mechanism (chem_opt=6) and MOSAIC using 8 sectional aerosol bins) including some aqueous reactions

For direct and indirect effect use: `phot_opt=2; ra_sw_physics=2; progn=1; mp_physics=2; aer_ra_feedback=1; wetscav_onoff=1; cldchem_onoff=1`

For dust and seasalt use `dust_opt=2, seas_opt=2`.

= 11 RADM2 chemical mechanism and MADE/SORGAM aerosols including some aqueous reactions

For direct and indirect effect use: `phot_opt=2; ra_sw_physics=2; progn=1; mp_physics=2; aer_ra_feedback=1; wetscav_onoff=1; cldchem_onoff=1`

For dust and seasalt use `dust_opt=2, seas_opt=2`.

KPP in WRF-Chem

Kinetic PreProcessor (KPP) reads chemical reactions and rate constants from ASCII input files and automatically generates code for chemistry integration using the Rosenbrock solver

No KPP for aerosols!


Advantages:

- ❑ less time consuming than manual coding
- ❑ less error prone
- ❑ numerically efficient
- ❑ flexibility in updating mechanism with additional species and equations
- ❑ suitable for adjoint code development

References:

- Damian, V., et al. (2002), The kinetic preprocessor KPP - a software environment for solving chemical kinetics, *Comput. Chem. Eng.*, 26(11), 1567-1579.
- Sandu, A., and R. Sander (2006), Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, *Atmos. Chem. Phys.*, 6, 187-195.
- Verwer, J., Spee, E., Blom, J. G., and Hunsdorfer, W. (1999), A second order Rosenbrock method applied to photochemical dispersion problems, *SIAM Journal on Scientific Computing*, 20, 1456–1480.
- www.mpch-mainz.mpg.de/~salzmann/my_home/sub/wkc.html

WKC (WRF-Chem/KPP Coupler) (\$WRF-CHEM/chem/KPP/util/wkc/)



*WRF-CHEM3.3 User's
guide, 2012*

Input files of KPP

***.spc** file

Definition of chemical species as variable or fixed value.

***.eqn** file

Writing chemical reactions in kpp format

***.kpp** file

Model description, computer language, precision, integrator (e.g. Rosenbrock solver) etc.

***.def** file

User defined functions

(also check \$WRF-CHEM/chem/KPP/kpp/kpp-2.1/util/ WRF_conform/ UserRateLaws.f90)

Pre-defined variables in WKC

	KPP equation file	Equation file units	Registry
Photolysis rate	J(Pj_no2)	s ⁻¹	ph_no2
Temperature	TEMP	K	t_phy
Third body concentration	C_M	(molecular moist air) cm ⁻³	Calculated from density
Water vapor concentration	C_H2O	Molecules cm ⁻³	Calculated from qvapor

WRF-CHEM3.6 User's guide, 2013

How to add a new KPP chemistry mechanism to WRF-Chem?

- 1) Add a new mechanism to `$WRF-CHEM/Registry/registry.chem`
- 2) Add new variables to the registry if necessary (new species, namelist options etc.)
- 3) Create subdirectories in `$WRF-CHEM/chem/KPP/` “mechanisms” and “inc”
- 4) Create new KPP files with new reactions, rates and species
- 5) Modify some `$WRF-CHEM/chem/*.F` (e.g. `chem_driver.F`) programs in order to include a new chemical mechanism
- 6) Modify `$WRF-CHEM/chem/convert_emiss.F` for new species
- 7) Compile a new WRF-CHEM code and run using updated emission files

KPP requires some UNIX tools - flex, yacc, and sed to be installed on your system before compiling the code !

setenv FLEX_LIB_DIR = /usr/local/lib

setenv WRF_KPP =1

Example: Adding a new chemistry mechanism to WRF-Chem for the state of the art SOA parameterization

In order to implement the new SOA mechanism (chem_opt=108 in V3.5) we need to modify the RACM_ESRSLORG gas chemistry:

- ☐ Separate MBO from internal alkenes - OLI species
- ☐ Add SESQ (biogenic VOC) reactions

Adding the new chemistry package and species to \$WRF-CHEM/Registry/registry.chem

```
state real sesq ikjftb chem 1 - irhusdf=(bdy_interp:dt) "sesq" "SESQ concentration" "ppmv"  
state real mbo ikjftb chem 1 - irhusdf=(bdy_interp:dt) "mbo" "MBO concentration" "ppmv"
```

The new package RACM_SOA_VBS_KPP chem_opt==108

Chem array:

G a s e s :
so2,sulf,no2,no,o3,hno3,h2o2,ald,hcho,op1,op2,paa,ora1,ora2,n2o5,no3,pan,hc3,hc5,hc8,eth,co,ete,
olt,oli,tol,xyl,aco3,tpan,hono,hno4,ket,gly,mgly,dcb,onit,csl,iso,co2,ch4,udd,hket,api,lim,dien,macr,hac
e, ishp,ison,mahp,mpan,nald, **sesq,mbo**, cvasoa1,cvasoa2,cvasoa3,cvasoa4,
cvbsoa1,cvbsoa2,cvbsoa3, cvbsoa4,ho,ho2,

Aerosols:

so4aj,so4ai,nh4aj,nh4ai,no3aj,no3ai, asoa1j,asoa1i,asoa2j,asoa2i,asoa3j,asoa3i,asoa4j,asoa4i,
bsoa1j,bsoa1i,bsoa2j,bsoa2i,bsoa3j,bsoa3i,bsoa4j,bsoa4i,
orgpaj,orgpai,ecj,eci,p25j,p25i,anthe,seas,soila,nu0,ac0,corn

All species within “chem” array are advected and mixed by WRF-CHEM!

Some species are part of the “misc” array (not advected)

\$WRF-CHEM/Registry/registry.chem:

non-transported radical species for the RACM mechanism

state	real	addt	ikj	misc	1	-	r	"addt"	"Radicals" "ppm"
state	real	addx	ikj	misc	1	-	r	"addx"	"Radicals" "ppm"
state	real	addc	ikj	misc	1	-	r	"addc"	"Radicals" "ppm"
state	real	etep	ikj	misc	1	-	r	"etep"	"Radicals" "ppm"
state	real	oltp	ikj	misc	1	-	r	"oltp"	"Radicals" "ppm"
state	real	olip	ikj	misc	1	-	r	"olip"	"Radicals" "ppm"
state	real	cslp	ikj	misc	1	-	r	"cslp"	"Radicals" "ppm"
state	real	limp	ikj	misc	1	-	r	"limp"	"Radicals" "ppm"
state	real	hc5p	ikj	misc	1	-	r	"hc5p"	"Radicals" "ppm"
state	real	hc8p	ikj	misc	1	-	r	"hc8p"	"Radicals" "ppm"
state	real	tolp	ikj	misc	1	-	r	"tolp"	"Radicals" "ppm"
state	real	xylp	ikj	misc	1	-	r	"xylp"	"Radicals" "ppm"
state	real	apip	ikj	misc	1	-	r	"apip"	"Radicals" "ppm"
state	real	isop	ikj	misc	1	-	r	"isop"	"Radicals" "ppm"
state	real	hc3p	ikj	misc	1	-	r	"hc3p"	"Radicals" "ppm"

.....

Make a new subdirectory in \$WRFV-CHEM/chem/KPP/mechanisms/

The name of this directory should be the same as the package name in the Registry without the “_kpp” suffix.

\$WRFV-CHEM/chem/KPP/mechanisms/**racm_soa_vbs**

Copy these files from chem/KPP/mechanisms/racm_esrlsorg to the new subdirectory and rename them except “atoms_red”:

atoms_red

racm_soa_vbs.def

racm_soa_vbs.eqn

racm_soa_vbs.kpp

racm_soa_vbs.spc

racm_soa_vbs_wrfkpp.equiv

racm_soa_vbs.def file

If necessary update equation sets and rate constants etc.

```
#include atoms_red
```

```
#include ./racm_soa_vbs.spc
```

```
#include ./racm_soa_vbs.eqn
```

```
#INLINE F90_RATES
```

```
REAL(KIND=dp) FUNCTION k46( TEMP, C_M )
```

```
    REAL(KIND=dp), INTENT(IN) :: temp, c_m
```

```
    REAL(KIND=dp) :: k0, k2, k3
```

```
    k0=2.4E-14_dp * EXP(460._dp/TEMP)
```

```
    k2=2.7E-17_dp * EXP(2199._dp/TEMP)
```

```
    k3=6.5E-34_dp * EXP(1335._dp/TEMP) * c_m
```

```
    k46=k0+k3/(1+k3/k2)
```

```
END FUNCTION k46
```

```
.....
```

racm_soa_vbs.spc file:

```
#DEFVAR
```

```
O3 =IGNORE ;
```

```
H2O2 =IGNORE ;
```

```
.....
```

```
SESQ =IGNORE ;
```

```
MBO =IGNORE ;
```

racm_soa_vbs.kpp file:

```
#MODEL racm_soa_vbs
```

```
#LANGUAGE Fortran90
```

```
#DOUBLE ON
```

```
#INTEGRATOR WRF_conform/rosenbrock
```

```
.....
```


racm_soa_vbs.eqn file:

#EQUATIONS {} ;

photolysis:

{001:J01} NO₂+hv=O₃P+NO : j(Pj_no2) ;

{002:J02} O₃+hv=O₁D{+O₂} : j(Pj_o31d) ;

{003:J03} O₃+hv=O₃P{+O₂} : j(Pj_o33p)

{004:J04} HONO+hv=HO+NO : j(Pj_hno2) ;

{005:J05} HNO₃+hv=HO+NO₂ : j(Pj_hno3) ;

.....

chemical reactions:

{024:001} O₃P+M{O₂}=O₃ : (C_M *6.00D-34*(TEMP/300.0)**(-2.4)) ;

{025:002} O₃P+O₃=M {2O₂} : ARR2(8.00D-12 , 2060.0_dp, TEMP) ;

{026:003} O₁D + M = O₃P : .78084*ARR2(2.15D-11 , -110.0_dp, TEMP) +
.20946*ARR2(3.30D-11 , -55.0_dp , TEMP) ;

{027:004} O₁D+H₂O=HO+HO : ARR2(1.63D-10 , -60.0_dp, TEMP) ;

{028:005} O₃+HO=HO₂{+O₂} : ARR2(1.70D-12 , 940.0_dp, TEMP) ;

Adding new reactions to racm_soa_vbs.eqn file

```
{245:222} SESQ+HO=0.36 KET+0.3 HCHO+0.05 ORA1+0.19 OLIP           : 2.52D-10 ;  
{246:223} SESQ+O3=0.51 HCHO+0.85 ALD+0.039 ORA1+0.23 KET+0.053 ORA2  
           +0.63 HO                                           : 5.60D-16 ;  
{247:224} SESQ+NO3=0.9 OLNN+0.10 OLND+0.9 MACR             : 2.20D-11 ;  
{248:225} MBO+HO=OLIP                                         : ARR2( 1.33D-11 , -500.0_dp, TEMP ) ;  
{249:226} MBO+NO3=0.11 OLNN+0.89 OLND                       : ARR2( 8.64D-13 , -450.0_dp, TEMP ) ;  
{250:227} MBO+O3=0.02 HCHO+0.99 ALD+0.16 KET+0.30 CO+0.011 H2O2  
           +0.14 ORA2+0.07 CH4+0.22 HO2+0.63 HO+0.23 MO2  
           +0.12 KETP+0.06 ETH+0.18 ETHP   :ARR2( 4.40D-15 , 845.0_dp,  
TEMP ) ;
```

Reference:

Papiez, M. R., et al. (2009), The impacts of reactive terpene emissions from plants on air quality in Las Vegas, Nevada, *Atmos. Environ.*, 43(27), 4109-4123

Update some \$WRF-CHEM/chem/ subroutines

In order to call necessary subroutines for the new chemical mechanism, e.g. SO₂->SO₄ conversion

chem_driver.F:

.....

so2so4_select: SELECT CASE(config_flags%chem_opt)

CASE (RADM2SORG,RADM2SORG_KPP,RACMSORG_KPP, **RACM_SOA_VBS_KPP)**

CALL wrf_debug(15,'gocart so2-so4 conversion')

call so2so4(chem,p_so2,p_sulf,p_h2o2,p_QC,T_PHY,MOIST, &

.....

Also the following subroutines in chem/

chemics_init.F
module_input_chem_data.F
mechanism_driver.F
cloudchem_driver.F
module_wetscav_driver.F
aerosol_driver.F
dry_dep_driver.F
emissions_driver.F
module_bioemi_megan2.F
module_add_emiss_burn.F
module_ftuv_driver.F
optical_driver.F
module_optical_averaging.F
...

Update chem/convert_emiss.F and generate new anthropogenic and biogenic emissions input (wrfchemi_* and wrfbiochemi* files)

```
.....  
#ifdef DM_PARALLEL  
    IF (wrf_dm_on_monitor()) THEN  
        READ(26,'(12E9.2)') dumc1(ids:ide-1,jds:jde-1)  
    ENDIF  
    DM_BCAST_MACRO(dumc1)  
#else  
    READ(26,'(12E9.2)') dumc1(ids:ide-1,jds:jde-1)  
#endif  
    grid%sebio_sesq(ips:ipe ,jps:jpe ) = dumc1(ips:ipe ,jps:jpe )  
ENDIF  
.....
```

After compiling WRF-CHEM with KPP option

\$WRF-CHEM/chem/KPP/mechanisms/racm_soa_vbs:

Makefile -> ../../util/Makefile_kpp

Makefile_racm_soa_vbs

atoms_red

racm_soa_vbs.def

racm_soa_vbs.eqn

racm_soa_vbs.kpp

racm_soa_vbs.spc

racm_soa_vbs_wrfkpp.equiv

racm_soa_vbs.map

racm_soa_vbs_Function.f90

racm_soa_vbs_Global.f90

racm_soa_vbs_Initialize.f90

racm_soa_vbs_Integrator.f90

racm_soa_vbs_Jacobian.f90

racm_soa_vbs_JacobianSP.f90

racm_soa_vbs_LinearAlgebra.f90

racm_soa_vbs_Main.f90

racm_soa_vbs_Model.f90

racm_soa_vbs_Monitor.f90

racm_soa_vbs_Parameters.f90

racm_soa_vbs_Precision.f90

racm_soa_vbs_Rates.f90

racm_soa_vbs_Update_Rconst.f90

racm_soa_vbs_Util.f90

racm_soa_vbs_mex_Fun.f90

racm_soa_vbs_mex_Jac_SP.f90

WKC generated racm_soa_vbs.map file

Variable species

1 = SULF (n) 32 = HKET (r) 63 = LIMP (r)
2 = CO2 (n) 33 = O3P (r) 64 = HC5P (r)
3 = ORA1 (n) 34 = PHO (r) 65 = HC8P (r)
4 = ORA2 (n) 35 = H2O2 (r) 66 = HCHO (r)
5 = CVASOA1 (n) 36 = ADDT (r) 67 = TOLP (r)
6 = CVASOA2 (r) 37 = ADDX (r) 68 = XYLP (r)
7 = CVASOA3 (r) 38 = ETE (r) 69 = OLIP (r)
8 = CVASOA4 (r) 39 = ADDC (r) 70 = ONIT (r)
9 = CVBSOA1 (n) 40 = PAA (r) 71 = DCB (r)
10 = CVBSOA2 (r) 41 = ISON (r) 72 = XO2 (r)
11 = CVBSOA3 (r) 42 = SESQ (r) 73 = OLI (r)

.....
18 = HC5 (r) 49 = MBO (r) 80 = HC3P (r)

Additional notes

- 1) Always run “clean -a” command after you change any of KPP files
- 2) When you add chemical species for a new chem_opt to registry.chem, place the gases between “so2 ... ho2” and if you add aerosols then place them after “so4aj...”
- 3) Only species within “chem” array (not “misc”!) are used to initialize from previous simulation data when chem_in_opt=1
- 4) Check if the added mechanisms work with pre-existing initial and boundary conditions, emissions, photolysis rates, aerosol modules, dry and wet deposition rates: e.g. check module_dep_simple.F
- 5) You can also simulate some species as passive tracers such as chem_opt=13-15 (prescribed emissions only), e.g. CO
- 6) chem_opt=16,17 (with modeled fluxes and prescribed emissions for CO2 and CH4)

The background of the slide features a light blue gradient with faint, semi-transparent chemical structures, including benzene rings and various functional groups. Scattered throughout are several translucent blue spheres of varying sizes, some of which have a bright white highlight, giving them a three-dimensional appearance. At the bottom of the slide, there are stylized, flowing white and light blue shapes that resemble liquid or smoke.

QUESTIONS ?

racm_soa_vbs_wrfkpp.equiv file:

! use this file for species that have different

! names in WRF and KPP

!

! Currently case sensitive

!

! left column	right column
----------------------	---------------------

! name in WRF	name in KPP
----------------------	--------------------

rpho	pho
-------------	------------